

Permian climate change recorded in palynomorph assemblages of Mozambique (Moatize Basin, eastern Tete Province)

Götz A., Hancox P., Lloyd A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Knowledge of Late Permian biodiversity patterns, following the end-Guadalupian crisis, is still in its infancy, since most recent studies have focused on the end-Permian biotic crisis. The palynological record of southern Africa, however, reveals major climatic changes during the Late Permian. Here we report new palynological data from eastern Tete Province of Mozambique, documenting the change from cool to warm temperate climates during the Lopingian. This prominent climate signal was also detected recently in other depositional environments elsewhere in southern Africa, and thus enables interregional temporal correlations.

<http://dx.doi.org/10.1515/acpa-2017-0001>

Keywords

Coal, Moatize Basin, Mozambique, Palaeoclimate, Palynology, Permian

References

- [1] BACKHOUSE J. 1991. Permian palynostratigraphy of the Collie Basin, Western Australia. *Rev. Palaeobot. Palynol.*, 67: 237-314.
- [2] BALME B.E. 1970. Palynology of Permian and Triassic strata in the Salt Range and Surghar Range, West Pakistan: 305-453. In: Kummel B. & Teichert C. (eds.), *Stratigraphic Boundary Problems: Permian and Triassic of West Pakistan*, University Press of Kansas, Kansas.
- [3] CAIRNCROSS B. 2001. An overview of the Permian (Karoo) coal deposits of southern Africa. *J. African Earth Sci.*, 33: 529-562.
- [4] CATUNEANU O., WOPFNER H., ERIKSSON P.G., CAIRNCROSS B., RUBIDGE B.S., SMITH R.M.H. & HANCOX P.J. 2005. The Karoo basins of southcentral Africa. *J. African Earth Sci.*, 43: 211-253.
- [5] COSTA M., CASTRO L., FERNANDES P., PEREIRA Z. & MARQUES J. 2016. Palinostratigrafia e maturação orgânica do Karoo Inferior nas sondagens ETA 15 e ETA 71 da Bacia de Moatize-Minjova, província de Tete (Moçambique). *Geonovas*, 29: 71-80.
- [6] FALCON R.M.S. 1975. Palynostratigraphy of the Lower Karoo sequence in the central Sebungwe District, Mid-Zambezi Basin, Rhodesia. *Palaeontol. Afric.*, 18: 1-29.
- [7] FALCON R.M.S. 1978. A palynological comparison of Karoo sediments on opposite sides of the Rhodesian Watershed with stratigraphic application. unpubl. PhD Thesis, University of the Witwatersrand, Johannesburg.
- [8] FALCON R.M.S., PINHEIRO H.J. & SHEPERD P. 1984a. The palynobiostratigraphy of the major coal seams in the Witbank Basin with lithostratigraphic, chronostratigraphic and palaeoclimatic implications. *Comuni. Serv. Geol. Portugal*, 70: 215-243.
- [9] FALCON R.M.S., LEMOS DE SOUSA M.J., PINHEIRO H.J. & MARQUES M.M. 1984b. Petrology and Palynology of Mozambique coals-Mucanha-Vúzi Region. *Comuni. Serv. Geol. Portugal*, 70: 321-338.

- [10] FERNANDES P., COGNÉ N., CHEW D.M., RODRIGUES B., SANTOS JORGE R.C.G., MARQUES M., JAMAL D. & VASCONCELOS L. 2015. The thermal history of the Karoo Moatize-Minjova Basin, Tete Province, Mozambique: An integrated vitrinite reflectance and apatite fission track thermochronology study. *J. African Earth Sci.*, 101: 55-72.
- [11] GASTALDO R.A., KAMO S.L., NEVELING J., GEISSMAN J.W., BAMFORD M. & LOOY C.V. 2015. Is the vertebrate-defined Permian-Triassic boundary in the Karoo Basin, South Africa, the terrestrial expression of the end-Permian marine event? *Geology*, 43: 1-5.
- [12] GÖTZ A.E. & RUCKWIED K. 2014. Palynological records of the Early Permian postglacial climate amelioration (Karoo Basin, South Africa). *Palaeobiodiv. Palaeoenviro.*, 94: 229-235.
- [13] HANKEL O. 1992. Late Permian to Early Triassic microfloral assemblages from the Maji ya Chumvi Formation, Kenya. *Rev. Palaeobot. Palynol.*, 72: 129-147.
- [14] HERMANN E., HOCHULI P.A., BUCHER H. & ROOHI G. 2012. Uppermost Permian to Middle Triassic palynology of the Salt Range and Surghar Range, Pakistan. *Rev. Palaeobot. Palynol.*, 169: 61-95.
- [15] JHA N. 2006. Permian palynology from India and Africa-A phytogeographic paradigm. *J. Palaeontol. Soc. India*, 51: 43-55.
- [16] LINDSTRÖM S. 1996. Late Permian palynology of Fossilryggen, Vestfjella, Dronning Maud Land, Antarctica. *Palynology*, 20: 15-48.
- [17] LINDSTRÖM S. & MCLOUGHLIN S. 2007. Synchronous palynofloristic extinction and recovery after the end-Permian event in the Prince Charles Mountains, Antarctica: implications for palynofloristic turnover across Gondwana. *Rev. Palaeobot. Palynol.*, 145: 89-122.
- [18] LLOYD A. 2015. Tete Coal Basin-What do we know? What do we still have to learn? Presentation delivered at the Fossil Fuel Foundation Solid fossil fuels in southern Africa and Madagascar, March 2015.
- [19] LOPES G., PEREIRA Z., FERNANDES P. & MARQUES J. 2014. Datação palinológica dos sedimentos glaciogénicos da Formação Tíltica de Vúzi, sondagem ETA 65, Bacia Carbonífera de Moatize-Minjova, Moçambique: resultados preliminares. *Comunicações Geológicas*, 101: 481-484.
- [20] MCKAY M.P., WEISLOGEL A.L., FILDANI A., BRUNT R.L., HODGSON D.M. & FLINT S.S. 2015. U-PB zircon tuff geochronology from the Karoo Basin, South Africa: implications of zircon recycling on stratigraphic age controls. *Int. Geol. Rev.*, 57: 393-410.
- [21] MORY A.J. & BACKHOUSE J. 1997. Permian Stratigraphy and Palynology of the Carnarvon Basin, Western Australia. *Geol. Sur. West. Australia*, 46: 1-101.
- [22] MUGABE J.A. 1999. Karoo deposits of Zambezi Graben-Moatize and Tete City Mozambique; Sedimentary facies distribution and palynological approach. unpubl. PhD Thesis, Utrecht University, Utrecht.
- [23] NETO J.N. 1976. Coal deposits of Mozambique: 642-647. In: Muir W.L.G. (ed.), *Coal Exploration, Proceedings of the 1st International Coal Symposium*, London.
- [24] NYAMBE I.A. & UTTING J. 1997. Stratigraphy and palynostratigraphy, Karoo Supergroup (Permian and Triassic), mid-Zambesi Valley, southern Zambia. *J. African Earth Sci.*, 24: 563-583.
- [25] PEREIRA Z., FERNANDES P., LOPES G., MARQUES J. & VASCONCELOS L. 2016. The Permian-Triassic transition in the Moatize-Minjova Basin, Karoo Supergroup, Mozambique: A palynological perspective. *Rev. Palaeobot. Palynol.*, 226: 1-19.
- [26] PRICE P.L. 1997. Permian to Jurassic palynostratigraphic nomenclature of the Bowen and Surat Basins: 137-178. In: Green P.M. (ed.), *The Surat and Bowen Basins, South-East Queensland*. Queensland Department of Mines and Energy.
- [27] RUBIDGE B.S., ERWIN D.H., RAMEZANI J., BOWRING S.A. & DE KLERK W.J. 2013. High-precision temporal calibration of late Permian vertebrate biostratigraphy: U-Pb constraints from the Karoo Supergroup, South Africa. *Geology*, 41(3): 363-366.
- [28] SCOTese C. 2016. A new global temperature curve for the Phanerozoic. GSA Annual Meeting Denver, Colorado, Abstracts with Programs 48(7): Paper No. 74-31; doi: 10.1130/abs/2016AM-287167.
- [29] SOUZA P.A. & MARQUES-TOIGO M. 2005. Progress on the palynostratigraphy of the Permian strata in Rio Grande do Sul State, Paraná basin, Brazil. *Anais da Academia Brasileira de Ciências*, 77: 353-365.
- [30] SRIVASTAVA S.C. & JHA N. 1990. Permian-Triassic palynofloral transition in Godavari Graben, Andhra Pradesh. *Palaeobotanist*, 38: 92-97.
- [31] SUN Y., JOACHIMSKI M.M., WIGNALL P.B., YAN C., CHEN Y., JIANG H., WANG L. & LAI X. 2012. Lethally hot temperatures during the early Triassic greenhouse. *Science*, 338: 366-370.
- [32] RUCKWIED K., GÖTZ A.E. & JONES P. 2014. Palynological records of the Permian Eccla Group (South Africa): Utilizing climatic icehouse-greenhouse signals for cross basin correlations. *Palaeogeogr., Palaeoclim., Palaeoecol.*, 413: 167-172.
- [33] VASCONCELOS L.S. 2009. Coal in Mozambique. *Proceedings of the 3rd Symposium on Gondwana Coals*, Porto Alegre, Brazil 2009.

- [34] VASCONCELOS L.S. 2012. Overview of the Mozambique Coal Deposits. Proceedings of the 34th International Geological Congress 2012, Australian Geosciences Council, Brisbane, Australia.
- [35] VEIZER J. & PROKOPH A. 2015. Temperatures and oxygen isotopic composition of Phanerozoic oceans. *Earth-Science Reviews*, 146: 92-104.
- [36] VIJAYA 2011. Palynostratigraphy of subsurface Upper Permian and Mesozoic succession, Rakshitpur area, Raniganj Coalfield, West Bengal. *Palaeoworld*, 20: 61-74.
- [37] WHEELER A. & GÖTZ A.E. 2017. Palynofacies as a tool for high-resolution palaeoenvironmental and palaeoclimatic reconstruction of Gondwanan postglacial coal deposits: No. 2 Coal Seam, Witbank Coalfield (South Africa). *Palaeobiodiv. Palaeoenvir.* [in press]. DOI: 10.1007/s12549-016-0248-x
- [38] WHEELER A. & GÖTZ A.E. 2016. Palynofacies patterns of the Highveld coal deposits (Karoo Basin, South Africa): Clues to reconstruction of palaeoenvironment and palaeoclimate. *Acta Palaeobot.*, 56(1): 3-15.
- [39] WRIGHT R.P. & ASKIN R.A. 1987. The Permian-Triassic boundary in the southern Morondava Basin of Madagascar as defined by plant microfossils: 157-166. In: McKenzie G.D. (ed.), *Gondwana Six: Stratigraphy, Sedimentology, and Paleontology*, Geophysical Monograph Series, Washington.